2011-2972 2011-08-16

Bedlam® Insecticide

INSECTICIDE

PRESSURIZED PRODUCT

To Kill Bed Bugs, Bed Bug Eggs.

For Indoor Use Only in Homes and Non-Food Areas of Restaurants, Schools, Nursing Homes, Warehouses, Offices, Apartments, Hotels, Motels, Kennels, and Hospitals.

COMMERCIAL

REGISTRATION NO. 30075

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

GUARANTEE:

| d-Phenothrin | 0.40% |
|--------------------------------------|-------|
| N-Octyl bicycloheptene dicarboximide | 1.60% |

BEDLAM® and MGK® - registered trademarks of McLaughlin Gormley King® Company

KEEP OUT OF REACH OF CHILDREN

Net Weight_____

Manufactured For:

McLAUGHLIN GORMLEY KING COMPANY 8810 Tenth Avenue North Minneapolis, MN 55427 U.S.A.

CAUTION – SKIN IRRITANT

(763) 544-0341



DIRECTIONS FOR USE

USE RESTRICTIONS: In hospitals and nursing homes, remove patients before treatment. Ventilate room for two hours before returning patients.

THIS PRODUCT IS NOT FOR USE ON HUMANS.

Do not use in commercial food/feed processing, preparation, food/feed storage or serving areas. In the home, all food/feed processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Exposed food/feed should be covered or removed. Remove pets, birds and cover fish aquariums before spraying.

SHAKE WELL BEFORE EACH USE. Only applicators should be in treatment areas during application. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry. Allow sprayed articles and surfaces to dry thoroughly before using or replacing.

<u>APPLICATION</u>: Spray in an inconspicuous area to test for possible staining or discoloration. Evaluate test area after drying. Hold container upright with nozzle aimed away from you. Depress valve and spray from a distance of 20 to 25 centimeters. Spray until damp. Most effective results are achieved when used as part of a treatment protocol that includes physical, environmental and other chemical pest control measures.

TO KILL BED BUGS AND BED BUG EGGS: Apply as a crack and crevice treatment on and around baseboards, floorboards, bed frames, wall hangings, headboards, furniture, door and window frames, millwork and walls. Apply as a spot treatment to carpet, mattresses, box springs, walls, furniture, floor and floor coverings, closets (remove clothing and other articles before treatment), and window treatment hardware such as curtain rods. Before treating mattresses and box springs, remove bedding. Treat areas that may harbor bed bugs or their eggs, such as tufts, folds and seams. Spray until damp. Then thoroughly clean and air-out mattresses and box springs.

<u>PRECAUTIONS</u>: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Do not induce vomiting. Avoid breathing vapours or spray mist. May cause skin irritation. Avoid contact with skin or eyes. Applicators must wear a long-sleeved shirt and long pants, socks, and shoes. In case of contact, flush with plenty of water. Wash with soap and warm water after use. Obtain medical attention if irritation persists. Avoid contamination of food or feedstuff. This product is toxic to aquatic organisms. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. Do not apply directly to food, water or feed supplements. CONTENTS UNDER PRESSURE.

<u>FIRST AID</u>: In case of contact with skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. In case of contact with eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Contains petroleum distillate - vomiting may cause aspiration pneumonia. Treat symptomatically.

<u>DISPOSAL</u>: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

STORAGE: Store in a cool, dry place. Do not store near heat or open flame.

<u>NOTICE TO USER</u>: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.