Syringe Label

GROUP	4A	INSECTICIDE
GROUP	4A	INSECTICIDE

Seclira Cockroach Gel Bait Reservoir

FOR INDOOR USE ONLY TO CONTROL GERMAN COCKROACHES AND ORIENTAL COCKROACHES

COMMERCIAL				
Insecticide				
Paste				
ACTIVE INGREDIENT: Dinotefuran,	0.50%			
REGISTRATION NO. 33424	PEST CONTROL PRODUCTS ACT			
READ THE LABEL AND BROCHURE BEFORE USING				
	RING LIFE OR PROPERTY INVOLVING THIS Y OR NIGHT 1-800-454-2673			
NET CONTER	NTS:			

BASF Canada Inc. 5025 Creekbank Road Building A, 2nd Floor Mississauga, Ontario L4W 0B6

Questions? Call 1-877-371-2273

Inner Box Label

GROUP 4A INSECTICIDE

Seclira Cockroach Gel Bait Reservoir

FOR INDOOR USE ONLY TO CONTROL GERMAN COCKROACHES AND ORIENTAL COCKROACHES

COMMERCIAL			
Insecticide			
Paste			
ACTIVE INGREDIENT: Dinotefuran,	0.50%		
REGISTRATION NO. 33424	PEST CONTROL PRODUCTS ACT		
READ THE LABEL AND BROCHURE BEFORE USING			
IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL DAY OR NIGHT 1-800-454-2673			
NET CONTEN	ITS:		

BASF Canada Inc. 5025 Creekbank Road Building A, 2nd Floor Mississauga, Ontario L4W 0B6

Questions? Call 1-877-371-2273

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND PETS. CAUTION. DO NOT apply to areas accessible to children and pets. DO NOT apply to surfaces that may come into contact with food/feed. DO NOT use on or contaminate food or feed. Exposed bait must be cleaned up immediately with a damp cloth or sponge. Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before re-use.

Wear long-sleeved shirt, long pants, chemical-resistant gloves plus shoes and socks during application, clean-up and repair.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15 - 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eyes. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 min. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

STORAGE

Store in a cool dry area away from heat or open flame and inaccessible to children and pets.

To prevent contamination, store this product away from food or feed.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. For information on disposal of unused, unwanted product, contact the manufacturer or

2022-0975 2022-03-04

the provincial regulatory agency. Contact the provincial agency in case of a spill and for clean-up of spills.

DIRECTIONS FOR USE

For indoor use only to control German cockroaches and Oriental cockroaches. See brochure inside for directions for use.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Brochure

GROUP 4A INSECTICIDE

Seclira Cockroach Gel Bait Reservoir

FOR INDOOR USE ONLY

Controls German and Oriental Cockroaches

COMMERCIAL

Insecticide

Paste

ACTIVE INGREDIENT:

REGISTRATION NO. 33424

PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL DAY OR NIGHT 1-800-454-2673

READ THE LABEL AND BROCHURE BEFORE USING

NET CONTENTS: 30 g

BASF Canada Inc. 5025 Creekbank Road Building A, 2nd Floor Mississauga, Ontario L4W 0B6

1-877-371-2273

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND PETS. CAUTION. DO NOT apply to areas accessible to children and pets. DO NOT apply to surfaces that may come into contact with food/feed. DO NOT use on or contaminate food or feed. Exposed bait must be cleaned up immediately with a damp cloth or sponge. Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before re-use.

Wear long-sleeved shirt, long pants, chemical-resistant gloves plus shoes and socks during application, clean-up and repair.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF EMERGENCY INVOLVING THIS PRODUCT, CALL 1-800-454-2673.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE

Store in a cool dry area away from heat or open flame and inaccessible to children and pets.

To prevent contamination, store this product away from food or feed.

DISPOSAL

Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the provincial agency in case of a spill and for clean-up of spills.

DIRECTIONS FOR USE

Users should treat a small, inconspicuous area of the surface first to ensure there are no adverse effects such as staining or discolouration before making a full-scale application to a surface. To prevent staining, exposed bait should be cleaned up immediately with a damp cloth, paper towel or sponge and then discarded in trash.

How to Apply

<u>Crack and Crevice Treatment:</u> Crack and crevice is an application directly into narrow openings on the surface of the structure. It does not include the treatment of exposed surfaces. Narrow openings typically occur at expansion joints, utility entry points and along baseboards and mouldings.

<u>Spot Treatment:</u> A spot application is an open, localized placement of bait in areas inaccessible to children and pets. Locations of treatment may include under sinks and under heavy appliances (e.g. fridge or stove).—Do not use spot treatments in food/feed areas of food/feed handling establishments.

<u>Void Treatment:</u> An application into inaccessible, enclosed empty spaces of a structure. For example, hollow walls and suspended ceilings.

Use Restrictions

- For indoor use only.
- DO NOT apply to areas accessible to children and pets.
- DO NOT apply bait to surfaces which may reach excessive temperatures. Examples
 include portions of stoves, ovens, grills, fume hoods, steam tables, toasters, fryers,
 dishwashers and hot water pipes.
- DO NOT place baits on surfaces that are openly exposed such as the junction between wall and floor.
- DO NOT place treatments in locations where routine cleaning might remove bait or transfer bait to food or food preparation surfaces.
- DO NOT apply to areas where food/feed products may become contaminated.
- For use as a crack and crevice void and spot treatment only.
- Keep out of sunlight or excess heat.
- DO NOT apply gel to areas that have been recently sprayed with insecticide.
- DO NOT spray insecticide products over gel placements as this may make gel repellent and reduce its effectiveness.

Where to Apply

For indoor use only in:

- Commercial, industrial and residential buildings including apartments, garages, homes, hospitals, hotels, motels, nursing homes, food/feed handling establishments[†], schools, supermarkets, and warehouses.
- Modes of Transportation (e.g., aircrafts, buses, boats, ships, trains, trailers and trucks)

[†]Food/feed handling establishments are places other than private residences in which food/feed is held, processed, prepared or served. This includes food/feed processing plants, food/feed storage areas, meat and poultry processing plants, restaurants and supermarkets.

- Non-food/feed areas include garbage rooms, restrooms, floor drains (to sewers), entries
 and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets
 and storage areas (after canning or bottling) and serving areas when no food/feed is
 present.
- Food/feed areas include areas for receiving, storing, packing (canning, bottling, wrapping, boxing) and preparing food/feed, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food/feed is exposed and facility is in operation) are also considered food/feed areas.

Controls: German and Oriental Cockroaches

Application Instructions

Place bait into areas of buildings that are inaccessible to children and pets. Bait that is in areas that may be accessible to children or pets must be cleaned up immediately with a damp cloth, paper towel or sponge and then discarded in trash.

Seclira Cockroach Gel Bait Reservoir is for use in controlling German and Oriental cockroaches indoors.

This product will not stick to surfaces that are dusty or greasy. Clean areas prior to application.

To apply bait, remove cap from tip of applicator tube. Use suitable equipment when applying into cracks and crevices, void spaces and other locations where German or Oriental cockroaches may be hiding. Replace cap on applicator tube when finished dispensing bait.

Areas to treat include: cracks and crevices; holes; pipe chases; under drain plates; in or under trash containers; hidden surfaces around sinks and storage areas; behind baseboards; around doors and windows; inside, behind and under cabinets, drawers and shelving; under and behind appliances such as stoves and refrigerators; in attics and crawl spaces; hollow spaces in floors, walls, ceilings, equipment legs; between equipment and floors and different elements of construction. Bait may also be applied to the undersides of furniture.

2022-0975 2022-03-04

Numerous small application spots are preferable to fewer larger application spots. Bait points should be made every 10-20 centimetres in areas of cockroach activity and suspected harborage. At each bait point, apply a 3-6 mm diameter spot of bait (equal to 0.25-0.50 g of bait) at the following rates:

- For light infestations of cockroaches, apply 4 6 bait points per 10 m². Light infestations would be indicated by sightings only at night when a light is turned on.
- For moderate infestations of cockroaches, apply 7-11 bait points per 10 m².
- For heavy infestations of cockroaches, apply 12 24 bait points per 10 m². Heavy infestations are indicated when cockroaches are readily seen upon the opening of cabinets, pantries or closets.

Re-apply when bait is consumed or is no longer palatable to the cockroaches. Applications should not be made more than once a month, unless bait is consumed. Use damp cloth, paper towel or sponge to remove any unwanted bait placements and discard in trash.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Seclira Cockroach Gel Bait Reservoir contains a Group 4A insecticide. Any insect population may contain individuals naturally resistant to Seclira Cockroach Gel Bait Reservoir and other Group 4 insecticides. The resistant individuals may dominate the insect population if these groups of insecticides are used repeatedly in the same locations. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed. To delay insecticide resistance:

- Where possible, rotate the use of Seclira Cockroach Gel Bait Reservoir or other Group 4
 insecticides with different groups that control the same pests in a location.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact BASF Canada Inc. at 1-877-371-2273.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.